

RESOURCES

- Download this card and additional resources at www.sprc.org or at www.stopasuicide.org
- Resource for implementing The Joint Commission 2007 Patient Safety Goals on Suicide www.sprc.org/library/jcsafetygoals.pdf
- SAFE-T drew upon the American Psychiatric Association Practice Guidelines for the Assessment and Treatment of Patients with Suicidal Behaviors www.psych.org/psych_pract/treatg/pg/SuicidalBehavior_05-15-06.pdf

ACKNOWLEDGEMENTS

- Originally conceived by Douglas Jacobs, MD, and developed as a collaboration between Screening for Mental Health, Inc. and the Suicide Prevention Resource Center.
- This material is based upon work supported by the Substance Abuse and Mental Health Services Administration (SAMHSA) under Grant No. 1U79SM57392. Any opinions/findings/conclusions/recommendations expressed in this material are those of the author and do not necessarily reflect the views of SAMHSA.

National Suicide Prevention Lifeline
1.800.273.TALK (8255)

COPYRIGHT 2007 BY EDUCATION DEVELOPMENT CENTER, INC., AND
SCREENING FOR MENTAL HEALTH, INC. ALL RIGHTS RESERVED.
PRINTED IN THE UNITED STATES OF AMERICA.
FOR NON-COMMERCIAL USE.



www.sprc.org



www.mentalhealthscreening.org

SAFE-T

Suicide Assessment Five-step Evaluation and Triage

1

IDENTIFY RISK FACTORS

Note those that can be modified to reduce risk

2

IDENTIFY PROTECTIVE FACTORS

Note those that can be enhanced

3

CONDUCT SUICIDE INQUIRY

Suicidal thoughts, plans behavior and intent

4

DETERMINE RISK LEVEL/INTERVENTION

Determine risk. Choose appropriate intervention to address and reduce risk

5

DOCUMENT

Assessment of risk, rationale, intervention and follow-up

National Suicide Prevention Lifeline
1.800.273.TALK (8255)

Suicide assessments should be conducted at first contact, with any subsequent suicidal behavior, increased ideation, or pertinent clinical change; for inpatients, prior to increasing privileges and at discharge.

1. RISK FACTORS

- ✓ **Current/past psychiatric diagnoses:** especially mood disorders, psychotic disorders, alcohol/substance abuse, Cluster B personality disorders. Co-morbidity and recent onset of illness increase risk
- ✓ **Key symptoms:** anhedonia, impulsivity, hopelessness, anxiety/panic, global insomnia, command hallucinations
- ✓ **Suicidal behavior:** history of prior suicide attempts, aborted suicide attempts or self-injurious behavior
- ✓ **Family history:** of suicide, attempts or Axis 1 psychiatric diagnoses requiring hospitalization
- ✓ **Precipitants/stressors:** triggering events leading to humiliation, shame or despair (i.e., loss of relationship, financial or health status—real or anticipated). Ongoing medical illness (esp. CNS disorders, pain). History of abuse or neglect. Intoxication
- ✓ **Access to firearms**

2. PROTECTIVE FACTORS *Protective factors, even if present, may not counteract significant acute risk*

- ✓ **Internal:** ability to cope with stress, religious beliefs, frustration tolerance, absence of psychosis
- ✓ **External:** responsibility to children or beloved pets, positive therapeutic relationships, social supports

3. SUICIDE INQUIRY *Specific questioning about thoughts, plans, behaviors, intent*

- ✓ **Ideation:** frequency, intensity, duration—in last 48 hours, past month and worst ever
 - ✓ **Plan:** timing, location, lethality, availability, preparatory acts
 - ✓ **Behaviors:** past attempts, aborted attempts, rehearsals (tying noose, loading gun), vs. non-suicidal self injurious actions
 - ✓ **Intent:** extent to which the patient (1) expects to carry out the plan and (2) believes the plan/act to be lethal vs. self-injurious; Explore ambivalence: reasons to die vs. reasons to live
- * Homicide Inquiry: when indicated, esp. postpartum, and in character disordered or paranoid males dealing with loss or humiliation. Inquire in four areas listed above.*

4. RISK LEVEL/INTERVENTION

- ✓ **Assessment of risk level** is based on clinical judgment, after completing steps 1–3
- ✓ **Reassess** as patient or environmental circumstances change

RISK LEVEL	RISK / PROTECTIVE FACTOR	SUICIDALITY	POSSIBLE INTERVENTIONS
High	Psychiatric diagnoses with severe symptoms, or acute precipitating event; protective factors not relevant	Potentially lethal suicide attempt or persistent ideation with strong intent or suicide rehearsal	Admission generally indicated unless a significant change reduces risk. Suicide precautions
Moderate	Multiple risk factors, few protective factors	Suicidal ideation with plan, but no intent or behavior	Admission may be necessary depending on risk factors. Develop crisis plan. Give emergency/crisis numbers
Low	Modifiable risk factors, strong protective factors	Thoughts of death, no plan, intent or behavior	Outpatient referral, symptom reduction. Give emergency/crisis numbers

(This chart is intended to represent a range of risk levels and interventions, not actual determinations.)

- ## 5. DOCUMENT:
- Risk level and rationale; treatment plan to address/reduce current risk (i.e., medication, setting, E.C.T., contact with significant others, consultation); firearm instructions, if relevant; follow up plan